



MEDLEY PHARMACEUTICALS

CIPROLEY CAPSULES

(Ciprofloxacin BP)

DESCRIPTION: CIPROLEY is a formulation of Ciprofloxacin HCl (BP) in the form of capsules.

COMPOSITION:

CIPROLEY 250mg CAPSULES:

Each capsule contains:

Ciprofloxacin HCl equivalent to Ciprofloxacin BP.....250mg

CIPROLEY 500mg CAPSULES:

Each capsule contains:

Ciprofloxacin HCl equivalent to Ciprofloxacin BP.....500mg

INTRODUCTION:

PHARMACODYNAMICS:

Inhibits DNA gyrase in susceptible organisms. Inhibits relaxation of super coiled DNA and promotes breakage of double stranded DNA.

PHARMACOKINETICS:

Well absorbed following the oral administration. Bioavailability is 70-80%. The maximum blood concentration is reached just 60-90mins after ingestion. The half life ($t_{1/2}$) is between 3-5hours. Ciprofloxacin is present in high concentration at the sites of infections e.g. in the body fluids and tissues.

INDICATIONS:

Treatment of documented or suspected infections of the lower respiratory tract, sinuses, skin/skin structure, bones/ joints and urinary tract including prostatitis. Due to susceptible bacterial strains, especially indicated for pseudomonas infections and those due to multi drug resistant Gram negative organisms. Chronic bacterial prostatitis, infectious diarrhea, complicated gram negative and anaerobic intra abdominal infections (with metronidazole) due to E.coli (Enteropathic strains), B.fragilis, P.marabilis, K.pneumonia, P. aeruginosa, Campylobacter jejuni or Shigella; approved for acute sinusitis caused by H.influenzae or M.catarrhalis, also used to treat typhoid fever due to Salmonella typhi (although eradication of the chronic typhoid carrier stage has not been proven), Osteomyelitis when parenteral therapy is not feasible, and sexually transmitted diseases such as uncomplicated cervical and urethral gonorrhoea due to Neisseria gonorrhoeae. Used ophthalmologically for superficial ocular infections, (Corneal ulcers, and conjunctivitis) due to susceptible strains.

CONTRAINDICATIONS:



Manufactured by:
ISO 9001-2008 Certified Company
Medley Pharmaceuticals
41-A, P.S.I.E. Jhang Bahtar Road,
Wah Cantt- Rawalpindi



MEDLEY PHARMACEUTICALS

Hypersensitivity to Ciprofloxacin and to other Quinolone chemotherapeutics.

DOSAGE:

Oral: 20-30mg /kg/day in 2 divided doses, maximum 1.5g/day.

UTI's: ADULTS: 250mg to 500mg every 12 hrs for 07 -10 Days following oral dose, depending on severity of infection and susceptibility.

LOWER RTI'S AND SKIN: 500-750mg twice daily for 7-14 days following oral dose and 400mg every 12 hrs for 7-14 days .

BONE/ JOINT INFECTIONS: 500-750mg twice daily for 4 to 6 weeks.

INFECTIOUS DIARRHEA: 500mg every 12 hrs for 5-7 days.

TYPHOID FEVER: 500mg every 12 hrs for 10 days.

URETHRAL/ CERVICAL GONOCOCCAL INFECTIONS: 250-500mg as a single dose. (Along with Doxycycline or azithromycin)

DISSEMINATED GONOCOCCAL INFECTIONS: 500mg twice daily to complete 7 days of therapy.

MILD TO MODERATE SINUSITIS: 500mg every 12 hrs for 10 days.

DRUG INTERACTIONS:

Should not be co-administered with iron, sucralfate or antacids containing magnesium, aluminium or calcium. Increased blood levels of theophylline are observed when administered along with Ciprofloxacin and increased creatinine levels are observed when ciprofloxacin is co-administered with Cyclosporin. The concomitant administration of Ciprofloxacin with Warfarin may intensify the action of warfarin. So serum levels should be monitored at regular interval when concomitant use is desirable. Metoclopramide increases the absorption of Ciprofloxacin causing less time to attain peak plasma concentration.

SIDE EFFECTS:

Nausea, Diarrhea, vomiting, dyspepsia, abdominal pain, flatulence and anorexia, dizziness, headache, tiredness, agitation, trembling, peripheral paresthesia, increased intracranial pressure, confusion, depression, hallucination, skin reaction, pruritus, tachycardia and very rarely hot flashes, migraine, fainting.

PRECAUTIONS:

Caution in the use of ciprofloxacin should be observed in elderly patients and in patients who have suffered previously from CNS disorders, i.e. previous history of convulsions, reduced cerebral blood flow, in epileptics, caution is necessary in patients having concomitant therapy with theophylline and in patients with impaired renal function .Patient should maintain an adequate fluid intake during treatment with ciprofloxacin.

PACK AND PRESENTATION:

CIPROLEY 250mg CAPSULES: Alu-PVC blister pack of 1x10's

CIPROLEY 500mg CAPSULES: Alu-PVC blister pack of 1x10's



Manufactured by:
ISO 9001-2008 Certified Company
Medley Pharmaceuticals
41-A, P.S.I.E. Jhang Bahtar Road,
Wah Cantt- Rawalpindi



MEDLEY PHARMACEUTICALS

STORAGE:

Keep in a cool and dry place.
Protect from light.
Keep out of the reach of children.



For further information please contact

Manufactured by:

ISO 9001-2008 Certified Company

Medley Pharmaceuticals

41-A, P.S.I.E Jhang Bahtar road,

Wah Cantt- Rawalpindi



Manufactured by:

ISO 9001-2008 Certified Company

Medley Pharmaceuticals

41-A, P.S.I.E. Jhang Bahtar Road,

Wah Cantt- Rawalpindi